


Orthopaedic and Sports Injuries Services "OASIS"
Munawar Shah FRCS, FRCS Tr & Orth
Consultant Trauma & Orthopaedic Surgeon

Little Aston Spire Hospital
Little Aston Hall Dr
Sutton Coldfield, B74 3UP
01215807406
01922656972
la.oasis@live.co.uk
<http://littleastonoasis.com>



Total Knee & Uni Knee replacement Exercise Guide

Regular exercise to restore your knee mobility and strength and a gradual return to everyday activities are important for your full recovery. Mr M Shah and physical therapist may recommend that you exercise approximately 20 to 30 minutes two or three times a day and walk 30 minutes, two or three times a day during your early recovery.

Mr M Shah may suggest some of the following exercises.

The following guide can help you better understand your exercise/activity program, supervised by your therapist and orthopaedic surgeon.

Early Postoperative Exercises

Start the following exercises as soon as you are able. You can begin these in the recovery room shortly after surgery. You may feel uncomfortable at first, but these exercises will speed your recovery and actually diminish your postoperative pain.

Quadriceps Sets

Tighten your thigh muscle. Try to straighten your knee. Hold for 5 to 10 seconds.

Repeat this exercise approximately 10 times during a two-minute period, rest one minute and repeat. Continue until your thigh feels fatigued.

Straight Leg Raises



Tighten the thigh muscle with your knee fully straightened on the bed, as with the Quad set. Lift your leg several inches. Hold for five to 10 seconds. Slowly lower.

Repeat until your thigh feels fatigued.

You also can do leg raises while sitting. Fully tighten your thigh muscle and hold your knee fully straightened with your leg unsupported. Repeat as above. Continue these exercises periodically until full strength returns to your thigh.


Ankle Pumps



Move your foot up and down rhythmically by contracting the calf and shin muscles. Perform this exercise periodically for two to three minutes, two or three times an hour in the recovery room.

Continue this exercise until you are fully recovered and all ankle and lower-leg swelling has subsided.


Knee Straightening Exercises



Place a small rolled towel just above your heel so that it is not touching the bed. Tighten your thigh. Try to fully straighten your knee and to touch the back of your knee to the bed. Hold fully straightened for five to 10 seconds.


Repeat until your thigh feels fatigued.

Bed-Supported Knee Bends




Bend your knee as much as possible while sliding your foot on the bed. Hold your knee in a maximally bent position for 5 to 10 seconds and then straighten.

Repeat several times until your leg feels fatigued or until you can completely bend your knee.



Ankle Pumps
flex ankle up and down




Quadriceps Sets:
tighten thigh muscles and hold for 5 seconds



Heel Slides:
Flex your hip and knee. Return knee to straight position.

Leg Lifts:
Raise leg six inches above the floor,
keeping knee straight.



Sitting Supported Knee Bends



While sitting at bedside or in a chair with your thigh supported, place your foot behind the heel of your operated knee for support. Slowly bend your knee as far as you can. Hold your knee in this position for 5 to 10 seconds.

Repeat several times until your leg feels fatigued or until you can completely bend your knee.

Sitting Unsupported Knee Bends




While sitting at bedside or in a chair with your thigh supported, bend your knee as far as you can until your foot rests on the floor. With your foot lightly resting on the floor, slide your upper body forward in the chair to increase your knee bend. Hold for 5 to 10 seconds. Straighten your knee fully.

Repeat several times until your leg feels fatigued or until you can completely bend your knee.

Early Activity

Soon after your surgery, you will begin to walk short distances in your hospital room and perform everyday activities. This early activity aids your recovery and helps your knee regain its strength and movement.

Walking



Proper walking is the best way to help your knee recover. At first, you will walk with a walker or crutches. Your surgeon or therapist will tell you how much weight to put on your leg.

Stand comfortably and erect with your weight evenly balanced on your walker or crutches. Advance your walker or crutches a short distance; then reach forward with your operated leg with your knee straightened so the heel of your foot touches the floor first. As you move forward, your knee and ankle will bend and your entire foot will rest evenly on the floor. As you complete the step, your toe will lift off the floor and your knee and hip will bend so that you can reach forward for your next step.


Remember, touch your heel first, then flatten your foot, and then lift your toes off the floor.



Walk as rhythmically and smooth as you can. Don't hurry. Adjust the length of your step and speed as necessary to walk with an even pattern. As your muscle strength and endurance improve, you may spend more time walking. You will gradually put more weight on your leg. You may use a cane in the hand opposite your surgery and eventually walk without an aid.

When you can walk and stand for more than 10 minutes and your knee is strong enough so that you are not carrying any weight on your walker or crutches (often about two to three weeks after your surgery), you can begin using a single crutch or cane. Hold the aid in the hand opposite the side of your surgery. You should not limp or lean away from your operated knee.

Stair Climbing and Descending



The ability to go up and down stairs requires strength and flexibility. At first, you will need a handrail for support and will be able to go only one step at a time. Always lead up the stairs with your good knee and down the stairs with your operated knee. Remember, "Up with the good" and "down with the bad." You may want to have someone help you until you have regained most of your strength and mobility.

Stair climbing is an excellent strengthening and endurance activity. Do not try to climb steps higher than the standard height (7 inches) and always use a handrail for balance. As you become stronger and more mobile, you can begin to climb stairs foot over foot.

Advanced Exercises and Activities

Once you have regained independence for short distances and a few steps, you may increase your activity. The pains of your knee problems before surgery and the pain and swelling after surgery have weakened your knee. A full recovery will take many months. The following exercises and activities will help you recover fully.

Standing Knee Bends


Standing erect with the aid of a walker or crutches, lift your thigh and bend your knee as much as you can. Hold for 5 to 10 seconds. Then straighten your knee, touching the floor with your heel first. Repeat several times until



Assisted Knee Bends

Lying on your back, place a folded towel over your operated knee and drop the towel to your foot. Bend your knee and apply gentle pressure through the towel to increase the bend.

Hold for 5 to 10 seconds; repeat several times until fatigued.



Knee Exercises with Resistance

You can place lightweights around your ankle and repeat any of the above exercises. These resistance exercises usually can begin four to six weeks after your surgery. Use one- to two-pound weights at first; gradually increase the weight as your strength returns. (Inexpensive wrap-around ankle weights with Velcro straps can be purchased at most sporting goods stores.)

Exercycling



Exercycling is an excellent activity to help you regain muscle strength and knee mobility. At first, adjust the seat height so that the bottom of your foot just touches the pedal with your knee almost straight. Peddle backward at first. Ride forward only after a comfortable cycling motion is possible backwards.


As you become stronger (at about four to six weeks) slowly increase the tension on the Exercycle. Exercycle for 10 to 15 minutes twice a day, gradually build up to 20 to 30 minutes, three or four times a week.

Pain or Swelling after Exercise You may experience knee pain or swelling after exercise or activity. You can relieve this by elevating your leg and applying ice wrapped in a towel. Exercise and activity should consistently improve your strength and mobility. If you have any questions or problems, contact your orthopaedic surgeon or physical therapist.

Orthopaedic and Sports Injuries Services "OASIS"	
Munawar Shah FRCS, FRCS Tr & Orth	
Consultant Trauma & Orthopaedic Surgeon	
<p>Little Aston Spire Hospital Little Aston Hall Dr Sutton Coldfield, B74 3UP 01215807406 01922656972 la.oasis@live.co.uk http://littleastonoasis.com</p>	A medical illustration of a human knee joint. It shows the femur (thigh bone) and tibia (shin bone) in a yellowish-brown color. Several surgical pins and wires are visible, indicating a surgical procedure or repair. The background is white.

Typical Home Exercises:


Instructions in your home exercise program may include some of the exercises shown below. Consult your therapist regarding the appropriateness of the exercises and the number of repetitions.




Ankle Pumps: Flex the ankle back and forth.




Quadriceps Sets: Tighten thigh muscles and hold for five seconds.




Heel slides: Flex your hip and knee, then return knee to the straight position



Leg lifts: Raise leg six inches above the floor, keeping knee straight.



Knee extensions: Place a pillow under your knee. Lift your foot off the mat.



Knee flexion stretch: Place a towel under your heel. Pull your knee toward your chest. Hold your knee in this flexed position for 15-20 seconds. Return knee to the straight position.